












Mollusks (*Gastropoda*) as Intermediate Hosts of Cattles' Trematodes (*Trematoda*) in Conditions of Dnipro Basin's Small Ponds (Northern Ukraine)

Diana Valeriyivna FESHCHENKO¹ , Tetiana Ivanivna BAKHUR^{2*} , Biricik Halil SELÇUK³ , Anatoliy Anatoliyovych ANTIPOV² , Oksana Anatoliyivna ZGHOZINSKA¹ , Oksana Anatoliyivna DUBOVA¹ , Valentina Oleksandrivna YEVSTAFYEVA⁴ , Volodymyr Petrovich GONCHARENKO² , Raisa Volodymyrivna SHAHANENKO² , Volodymyr Sergiyovych SHAHANENKO² , Vitaliy Vasyliovych MELNYCHUK⁴ 

¹Department of Parasitology, Veterinary-Sanitary Expertise and Zoohygiene, Zhytomyr National Agroecological University, Zhytomyr, Ukraine

²Department of Parasitology and Pharmacology, Bila Tserkva National Agrarian University, Bila Tserkva, Kiev, Ukraine

³Department of Laboratory Animals, Afyon Kocatepe Üniversitesi, Afyon, Turkey

⁴Department of Parasitology and Veterinary-Sanitary Expertise, Poltava State Agrarian Academy, Poltava, Ukraine

Cite this article as: Feshchenko, D.V., Bakhur, T.I., Biricik, H.S., Antipov, A.A., Zghozinska, O.A., Dubova, O.A., Yevstafyeva, V.O., Goncharenko, V.P., Shahanenko, R.V., Shahanenko, V.S., Melnychuk, V.V., 2019. Mollusks (*Gastropoda*) as Intermediate Hosts of Cattles' Trematodes (*Trematoda*) in Conditions of Dnipro Basin's Small Ponds (Northern Ukraine). *Acta Vet Eurasia* 2019; 45: 16-21.

ORCID IDs of the authors: D.V.F. 0000-0002-4811-2488; T.I.B. 0000-0001-8271-8267; B.H.S. 0000-0003-4974-1611; A.A.A. 0000-0003-3955-3377; O.A.Z. 0000-0003-4622-6307; O.A.D. 0000-0002-8303-573X; V.O.Y. 0000-0003-4809-2584; V.P.G. 0000-0002-7279-6146; R.V.S. 0000-0002-5848-1367; V.S.S. 0000-0002-3484-2962; V.V.M. 0000-0003-1927-1065.

Abstract

The article presents the data on distribution and defeat of gastropod mollusks by parasitic trematodes in biotopes of small reservoirs (rivers, lakes and swamps) of the Dnipro basin of northern regions of Ukraine. During the 2016-2017 years, at the following areas were collected and identified: *Lymnaea* (*L. stagnalis* (Linnaeus, 1758); *L. truncatula* (Müller, 1774); *Planorbis* (*P. corneus* (Linnaeus, 1758); *P. planorbis* (Linnaeus, 1758); *Viviparus contectus* (Millet, 1813); *Valvata piscinalis* (Müller, 1774) and *Succinea pfeifferi* (Rossmässler, 1834). The microscopic study of the mollusks' liver allowed us to detect the presence of pathogens of cattle trematodoses inside a cer-

tain number of the snails – *Fasciola hepatica* (Linnaeus, 1758) and *Paramphistomum* sp. (Fischoeder, 1901). At biotopes of small rivers, 8.3% of mollusks *L. truncatula* species, 23.5% of *L. stagnalis* and 5.7% of *P. corneus* were affected. At lakes and swamps, the number of affected *L. truncatula* was 36.3%, and *L. stagnalis* – 13.7%. It was determined the defeat of ruminants with fasciolosis and paramphistomatoses in designated regions. It testifies to the formation of sustainable natural foci of these invasions.

Keywords: Cattle, *Fasciola hepatica*, Gastropod mollusks, *Paramphistomum* sp., trematodes

Introduction

The parasitic system is a special biosystem based on the trophic relationship between the parasite and the host (Beklemishev, 1970). The functioning of parasitic systems, in particular, the regulation of the relationship between their members, is ensured by a whole complex of environmental factors (Antipov et al., 2018; Kennedy, 1978; Krasnoshchekov, 1996).

Trematodes (*Trematoda*: *Digenea*) are extremely common in tropical and subtropical climates (including Bolivia, Ecuador, Peru, Cuba, Egypt, Turkey, Iran), as well as in the European Union, Russia and Ukraine. Intermediate hosts of trematodes are freshwater gastropods (Caminade et al., 2015; Cañete et al., 2004; Caron et al., 2014; El-Shazly et al., 2012; Gorokhov et al., 2010; Khoramian et al., 2014; Lopez et al., 2012; Novobilský et al., 2013; Rondelaud et al., 2015; Stadnichenko, 2006).

Trematodes, which parasitize in the organs of the digestive system of cattle (in particular in the duodenum, the liver and the sections of the multi-chamber stomach) are an important epizootic danger to the cattle breeding of these countries (Munguía-Xóchihua et al., 2007). In this context, separate representatives of the following species should be singled out: *Fasciola hepatica* (*F. hepatica*) (Linnaeus, 1758), *Paramphistomum ichikawai* (Fukui, 1922), *Paramphistomum cervi* (Schränk, 1790) and *Liorchis scotiae* (Willmott, 1950).

Gastropods mollusks (*Gastropoda*) the most numerous class in the *Mollusca* type (Linnaeus, 1758), which has about 60.000-75.000 species (Zhadin, 1926). Among the gastropods, there are very few real parasites, for example: *Eulima bilineata* (Alder, 1848), parasite of bottom marine animals (*Ophiuroidea*; Grey, 1840) from the type of echinoderms (*Echinodermata*; Klein, 1734) (Nekhaev, 2011).

Address for Correspondence: Tetiana Ivanivna BAKHUR • E-mail: fly_13@ukr.net

Received Date: 6 September 2018 • Accepted Date: 14 February 2019 • DOI: 10.26650/actavet.2019.18009

© Copyright 2019 by Official Acta Veterinaria Eurasia. Available online at actaveteurasia.istanbul.edu.tr



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

However, many species of gastropods are included to the life cycle of helminths and the epizootic chain of domestic animals' invasive diseases, in particular *Fasciola* and *Paramphistomum* invasion of cattle.

According to their biology, freshwater mollusks are natural inhabitants of large and small ponds involved in water purification and are indicator species for determining the degree of anthropogenic loading of biotopes. The fauna of mollusks depends to a large extent on the propagation of certain types of higher vegetation and depth of water bodies, velocity, temperature and pH of water in the pond (Bennema et al., 2011; Sargison et al., 2016; Zhytova and Korniyushyn, 2017).

In Ukraine, the northern Polissya regions, geographical center of which is in the territory of Kyiv and Zhytomyr regions, are characterized by the largest species diversity of gastropods in the small rivers of the Dnipro basin (Zhytova and Korniyushyn, 2017). It causes a severe enzootic situation with *Trematoda* invasions of large and small ruminants in this zone.



Figure 1. Points of mollusks' collecting at northern regions of Ukraine (Dnipro basin)



Figure 2. Shells of *Lymnaea truncatula* (Müller, 1774)

So, the study of the epizootology of ruminant animals' trematodes and freshwater fauna is closely interrelated and the provision of quality products and profitability of cattle breeding in endemic areas is impossible without comprehensive knowledge about ecology of the gastropods.

Materials and Methods

The research protocol of the current study was approved by the Ethic Committee of the Zhytomyr National Agroecological University (Approval number: 2016/07).

Freshwater mollusks were collected during 2016-2017 from river basins and their tributaries-rivers Teteriv and Sluch (Zhytomyr region), Bucha, Ros, Skvira (Kiev region). Particular attention was paid to the study of the malakofauna of lakes and swamps, which are located on the territory of these regions. Points of mollusks' collecting are shown on the map (Figure 1).

In general, more than one thousand specimens of mollusks from the family *Lymnaeidae* (2 species), *Valvatidae* (1 species), *Planorbidae* (2 species), *Viviparidae* (1 species) and *Succineidae* (1 species) were investigated. Cameral processing of materials was carried out in accordance with the recommendations of Zdun (1961). The mollusks were harvested using common methods (Stadnichenko, 2006). Identification of the species of mollusks was carried out according to external conchological features (Stadnichenko, 2004).

For histological study of the most common species of mollusks – *Lymnaea stagnalis* (*L. stagnalis*) (Linné, 1758), *Planorbarius corneus* (*P. corneus*) (Linné, 1758), hepatopancreas was taken from which were made preparations for parasitological microscopic examination. The presence of partenites (rediae) and larvae (cercariae) in the body of mollusks was determined using a microscope "XS-6320 (MICROmed, Poltava, Ukraine)". The intensity of the invasion was assessed visually according to the following criteria: weak-larvae defeat less than 1/10 volume of hepatopancreas; average-from 1/10 to 1/2; and strong – more than 1/2.

Results and Discussion

From the biotopes of groundwater ponds of the Dnipro basin in Kiev and Zhytomyr regions, we collected and identified 6 species of freshwater mollusks: *L. stagnalis* (Linnaeus, 1758); *L. truncatula* (Müller, 1774); *P. corneus* (Linnaeus, 1758); *P. planorbis* (Linnaeus, 1758); *Viviparus contectus* (Millet, 1813); *Valvata* (*V.*) *piscinalis* (Müller, 1774), and 1 terrestrial species – *Succinea* (*S.*) *pfeifferi* (Rossmässler, 1834).

Dominant species were *L. truncatula*, *L. stagnalis*, and *P. corneus* with a population density of 4-9 specimens/m² in the spring, in the summer of 14-20 specimens/m², and in the autumn, 1-3 specimens/m². Other species occurred singly, 1-3 specimens/m². For trematodes (*Trematoda: Digenea*) of ruminants, as intermediate hosts can serve: *L. truncatula* – for *Fasciola hepatica*

(Figure 2); *L. stagnalis* (Figure 3), *L. truncatula* and *P. corneus* (Figure 4) – for pathogens *Paramphistomidae* sp.; *S. pfeifferi* (Figure 5), sometimes may be an intermediate host of the dicrocoeliosis' pathogen (*Dicrocoelium lanceatum*) (Charlier et al., 2016; Dreyfuss et al., 1999; Faltýnková et al., 2008; Hodasi, 1972; Vignoles et al., 2016; Zhytova and Korniyushyn 2012).



Figure 3. Shells of *Lymnaea stagnalis* (Linnaeus, 1758)



Figure 4. Shells of *Planorbis corneus* (Linnaeus, 1758)



Figure 5. Shells of *Succinea pfeifferi* (Rossmässler, 1834)

Mollusk *V. piscinalis* can be the first intermediate host for trematodes *Ichthyocotylurus pileatus* (Rudolphi, 1802) and *Diplostomum baeri* (Dubois, 1937), the additional hosts of which are freshwater fish, and the definitive – gulls. According to the fact that there is no literature data about the distribution of ruminant trematodes' larvae in *V. piscinalis*, microscopic examination of these mollusks' hepatopancreas was not carried out. Our studies have confirmed that the ratio of species of molluscs in a certain area and the level of their invasion by parthenites and larvae of the trematodes depend on the biotope's characteristics.

Thus, the malakofauna of small flowing (rivers, their tributaries and floodplains) and standing reservoirs (lakes, swamps) of the Dnipro basin of northern Ukraine differed considerably.

The data obtained is illustrated in Table 1, from which it is evident that part of the representatives of the species *L. truncatula* (up to 36.3%) was affected by rediae and cercariae of *F. hepatica* (Figure 6); up to 23.5% of *L. stagnalis* mollusks contained parthenogenetic genera of both *F. hepatica* and *Paramphistomum* sp. in hepatopancreas; some specimens of gastropods of the




Figure 6. a, b. Parasitic trematode larvae (Trematoda: Digenea) in the liver of mollusk *Lymnaea truncatula* (Muller, 1774): a) mature cercariae; b) young cercariae




Figure 7. Average extensiveness of the invasion of large and small cattle by trematodes in the territory of Kyiv and Zhytomyr regions during 2016-2017, %

Table 1. Parasitologic study of mollusks' hepatopancreas (selected from ponds, Dnipro basin, Northern Ukraine)

Species of Mollusks	Extensiveness of the Invasion, %		Level of the Invasion	
	Biotores of Lakes and Swamps	Biotores of Rivers	Biotores of Lakes and Swamps	Biotores of Rivers
<i>L. truncatula</i> (Müller, 1774)			<i>F. hepatica</i>	
	36.3	8.3	Average	Weak
<i>L. stagnalis</i> (Linnaeus, 1758)			<i>F. hepatica</i> and <i>Paramphistomidae</i> sp.	
	13.7	23.5	Weak	Strong
<i>P. corneus</i> (Linnaeus, 1758)			<i>Paramphistomidae</i> sp.	
	0	5.7	0	Weak
<i>P. planorbis</i> (Linnaeus, 1758)			<i>Paramphistomidae</i> sp.	
	0	0	0	0
<i>S. pfeifferi</i> (Rossmässler, 1834)			<i>Dicrocoelium lanceatum</i>	
	0	0	0	0

species *P. corneus* (5.7%) were invasive solely by *Paramphistomum* sp. trematodes.

Taking into account that *L. truncatula* mollusks are biological hosts exclusively for *F. hepatica*, as well as the fact that they are widely distributed in the ponds of Ukrainian Polissya and the damage by the trematodes' partenites and larvae (especially in biotores of lakes and swamps), there is a high risk for ruminants about fasciolosis throughout the studied area.

In contrast, the populations of *L. stagnalis* species of mollusks, which are often infected with rediae and cercariae of *F. hepatica* and *Paramphistomidae* sp., are more numerous near rivers than near standing water bodies.

Thus, in the northern regions of Ukraine at ponds of Dnipro basin, the predominant number of cases of cattle contamination by pathogens of paramphistomoses occurs near the rivers, and invasion by fascioles – near any ponds (both flowing and standing).

To establish the scale of the damage by the ruminant animals' trematodes in the investigated area, we have analyzed the official data of the state veterinary service in the Kyiv and Zhytomyr regions for 2016-2017. The analysis results are presented in Figure 7. The high extensiveness of ruminal invasion with *Paramphistomidae* sp. (1.2-1.9% of all examined animals), and especially *F. hepatica* (3.6-10.8%), confirm the obtained data on the widespread distribution of the diseases in the northern regions of Ukraine.

It is known that 25% invasion of malakofauna with helminths in a particular region is quite sufficient for the preservation and spread of the disease among the favorable domestic animals (Beesley et al., 2017). Just in one mollusc can develop at the same time more than 100 trematodes' cercariae. This is confirmed by our own research, according to which, up to 11% of

cattle in the territory of Kyiv and Zhytomyr regions are having fasciolosis. It is not a critical indicator, but it is also not a reason to ignore the problem, since the stationary troubles of farms bring not less economic damage than sudden outbreaks of especially dangerous diseases.

To disrupt the stability of the parasitic system (trematodes-mollusks-ruminants), a sharp change in external environmental factors may occur. For example, the sublethal for gastropods is the temperature of water above 27°C, depending on the type of mollusk (Afanasyev, 1993), due to the population of mollusks in the waters of Central Europe in the summer, it is sometimes significantly reduced. However, representatives of the subclass *Pulmonata* (Cuvier, 1817), adapted to the sharp fluctuations of temperature, falling into the summer (aestivation period) or winter (hibernation period) lethargy.

In most of the water systems of the Dnipro basin in the north of Ukraine, conditions for the development of freshwater mollusks are quite favorable. The climate of this area is mild, moderately continental. Endemic species of molluscs are evolutionally adapted to possible short-term critical climatic oscillations, as evidenced by the seasonal fluctuations of the number of gastropods in the studied ponds.

Thus, the presence of a permanent population of gastropod mollusks, affected by trematodes' pathogens, in small ponds of the north of Ukraine creates a permanent natural reservoir of fasciolosis and paramphistomatoses. This does not let to allow the cattle to be protected from further invasion without using of complex control measures against helminths.

Conclusion

In the biotope of natural ponds of the Dnipro basin at the northern regions of Ukraine, there are 7 species of gastropods, which are intermediates of ruminant's trematodes: *L. stagnalis*,

L. truncatula, *P. corneus*, *P. planorbis*, *Viviparus contectus*, *V. piscinalis* and *S. pfeifferi*. The first three species are dominated by numbers: from 4 to 9 specimens/m² in spring, from 14 to 20 per m² in summer, and from 1 to 3 per m² in autumn.

Intraspecies indices of mollusks' invasiveness by partenites and larvae of the trematodes varied depending on the type of ponds. More often (23.5%) and more intensively the *L. stagnalis* population was affected by rediae and cercariae of *F. hepatica* and *Paramphistomum* sp., while in *P. corneus* (5.7%) was found only *Paramphistomum* sp. in rivers' biotopes. Indices of damage to mollusks of the species *L. truncatula* by the *F. hepatica* trematodes increased in lakes and swamps (36.3%).

Ethics Committee Approval: Ethics Committee approval was received for this study from the Ethics Committee of Zhytomyr National Agroecological University, Ukraine (Approval number: 2016/07).

Peer-review: Externally peer-reviewed.

Author Contributions: Concept – D.V.F., T.I.B., A.A.A.; Design – D.V.F., O.A.Z., R.V.S.; Supervision – D.V.F., A.A.A., V.P.G.; Resources – V.O.Y.; Materials – D.V.F., O.A.D., V.V.M.; Data Collection and/or Processing – T.I.B., V.P.G., V.S.S.; Analysis and/or Interpretation – D.V.F., T.I.B., B.H.S.; Literature Search – D.V.F., O.A.Z., O.A.D., V.S.S.; Writing Manuscript – D.V.F., T.I.B.; Critical Review – A.A.A., B.H.S.

Acknowledgements: The authors are grateful to Associate Professor Zhitova O.P. from Zhytomyr National Agroecological University for thematic advisory support.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

References

- Afanasyev, S.A., 1993.** Influence of elevated water temperatures on elimination of different-sized *Dreissenia*. In: Materials of the VI All-Union Conference "The species and its productivity in the range", Minsk, 115.
- Antipov, A.A., Bakhur, T.I., Feshchenko, D.V., Romanishina, T.A., Avramenko, N.V., Goncharenko, V.P., Zghozinska, O.A., Solovyova, L.M., Koziy, N.V., Pidborska, R.V., Shahanenko, V.S., Dzhmil, V.I., Tyshkivska, N.V., 2018.** Earthworms (*Lumbricidae*) as intermediate hosts of lung nematodes (*Metastrongylidae*) of swine in Kyiv and Zhytomyr regions of Ukraine. *Vestnik Zoologii* 52, 59-64.
- Beesley, N.J., Williams, D.J.L., Paterson, S., Hodgkinson, J., 2017.** *Fasciola hepatica* demonstrates high levels of genetic diversity, a lack of population structure and high gene flow: Possible implications for drug resistance. *International Journal for Parasitology* 47, 11-20.
- Beklemishev, V.N., 1970.** Disease Agents as Components of Biocenoses. In: *Biocenotic Basics of Comparative Parasitology*. Moscow, pp. 334-352.
- Bennema, S.C., Ducheyne, E., Vercruyse, J., Claerebout, E., Hendrickx, G., Charlier, J., 2011.** Relative importance of management, meteorological and environmental factors in the spatial distribution of *Fasciola hepatica* in dairy cattle in a temperate climate zone. *International Journal for Parasitology* 41, 225-233.
- Caminade, C., Van Dijk, J., Baylis, M., Williams, D., 2015.** Modelling recent and future climatic suitability for fasciolosis in Europe. *Geospatial Health* 9, 301-308.
- Cañete, R., Yong, M., Sánchez, J., Wong, L., Gutiérrez, A., 2004.** Population dynamics of intermediate snail hosts of *Fasciola hepatica* and some environmental factors in San Juan y Martínez municipality, Cuba. *Memórias do Instituto Oswaldo Cruz* 99, 257-262.
- Caron, Y., Martens, K., Lempereur, L., Saegerman, C., Losson, B., 2014.** New insight in lymnaeid snails (*Mollusca, Gastropoda*) as intermediate hosts of *Fasciola hepatica* (*Trematoda, Digenea*) in Belgium and Luxembourg. *Parasites and Vectors* 7, 66.
- Charlier, J., Ghebretinsae, A.H., Levecke, B., Ducheyne, E., Claerebout, E., Vercruyse, J., 2016.** Climate-driven longitudinal trends in pasture-borne helminth infections of dairy cattle. *International Journal for Parasitology* 46, 881-888.
- Gorokhov, V.V., Skira, V.N., Klenova, I.F., Taiichinov, U.G., Volichev, A.N., Peshkov, R.A., Maisheva, M.A., Gorokhova, E.V., Melnikova, L.E.E., Simolovskaya, N.A., Ermakov, I.V., 2010.** Epizootic situation in the main helminthiases in the Russian Federation. *Theory and Practice of Parasitic Animal Diseases* 11, 124-131.
- Dreyfuss, G., Vignoles, P., Rondelaud, D., Vareille-Morel, C., 1999.** *Fasciola hepatica*: characteristics of infection in *Lymnaea truncatula* in relation to the number of Miracidia at exposure. *Experimental Parasitology* 92, 19-23.
- El-Shazly, A.M., Nabih, N., Salem, D.A.B., Mohamed, M.Z., 2012.** Snail populations in Dakahlia Governorate, Egypt, with special reference to lymnaeids. *Egypt Journal of Biology* 14, 45-49.
- Faltýnková, A., Nasincová, V., Kablášková, L., 2008.** Larval trematodes (*Digenea*) of planorbisid snails (*Gastropoda: Pulmonata*) in Central Europe: a survey of species and key to their identification. *Systematic Parasitology* 69, 155-178.
- Hodasi, J.K.M., 1972.** The output of cercariae of *Fasciola hepatica* by *Lymnaea truncatula* and the distribution of metacercariae on grass. *Parasitology* 64, 53.
- Khoramian, H., Arbabi, M., Osqoi, M. M., Delavari, M., Hooshyar, H., Asgari, M., 2014.** Prevalence of ruminants fascioliasis and their economic effects in Kashan, center of Iran. *Asian Pacific Journal of Tropical Biomedicine* 4, 918-922.
- Kennedy, K., 1978.** *Ecological parasitology*. Moscow, Mir, p. 230.
- Krasnoshchekov, G.P., 1996.** Parasitic system: habitat and adaptation of parasites. *Tolyatti*, 50.
- Lopez, M., White, A.C.Jr., Cabada, M.M., 2012.** Burden of *Fasciola hepatica* Infection among Children from Paucartambo in Cusco, Peru. *The American Journal of Tropical Medicine and Hygiene* 86, 481-485.
- Munguía-Xóchihua, J.A., Ibarra-Velarde, F., Ducoing-Watty, A., Montenegro-Cristino, N., Quiroz-Romero, H., 2007.** Prevalence of *Fasciola hepatica* (ELISA and fecal analysis) in ruminants from a semi-desert area in the northwest of Mexico. *Parasitology Research* 101, 127-130.

- Nekhaev, I.O., 2011.** Two species of parasitic molluscs new for Russian seas. *Ruthenica* 21, 69-72.
- Novobilský, A., Kašný, M., Beran, L., Rondelaud, D., Höglund, J., 2013.** *Lymnaea palustris* and *Lymnaea fuscus* are potential but uncommon intermediate hosts of *Fasciola hepatica* in Sweden. *Parasites and Vectors* 6, 251.
- Rondelaud, D., Vignoles, P., Dreyfuss, G., 2015.** Larval trematode infections in *Galba truncatula* (*Gastropoda, Lymnaeidae*) from the Brenne Regional Natural Park, central France. *Journal of Helminthology* 90, 256-261.
- Sargison, N., Francis, E., Davison, C., Bronsvoort, B.M.C., Handel, I., Mazeri, S., 2016.** Observations on the biology, epidemiology and economic relevance of rumen flukes (*Paramphistomidae*) in cattle kept in a temperate environment. *Veterinary Parasitology* 219, 7-16.
- Stadnichenko, A.P., 2004.** Prothorovic and calyx (*Lymnaeidae, Acroloxidae*) of Ukraine. Kiev, Center for Educational Literature, 327.
- Stadnichenko, A.P., 2006.** *Lymnaeidae* and *Acroloxidae* of Ukraine: Methods of collection and study, biology, ecology, useful and harmful meaning: monograph. Zhytomyr, Ruta, 168.
- Vignoles, P., Rondelaud, D., Dreyfuss, G., 2016.** Aptitude of *Lymnaea palustris* and *L. stagnalis* to *Fasciola hepatica* larval development through the infection of several successive generations of 4-mm-high snails. *Parasitology Research* 115, 2263-2268.
- Zdun, V.I., 1961.** Larvae of Trematodes in Freshwater Molluscs of Ukraine. Ukrainian Academy of Sciences Press, Kiev, p. 141.
- Zhadin, V.I., 1926.** Biology of fresh-water mollusks of ephemeral bodies of water. *Russische Hydrobiologische Zeitschrift*, 131.
- Zhytova, E.P., Kornyushyn, V.V., 2012.** The dynamics of the dimensional and age structure of *Lymnaea (L.) stagnalis* (*Gastropoda, Pulmonata*) populations infected with trematode parthenitae and larvae. *Visnyk of Lviv University, Biological series* 58, 193-201.
- Zhytova, E.P., Kornyushyn, V.V., 2017.** The role of different mollusk species in maintaining the transmission of polyhostal *Trematode species* in Ukrainian Polissya waters: the specificity of *Trematode* parthenogenetic generations to mollusk hosts. *Vestnik Zoologii* 51, 295-310.